

# Local mirror symmetry and Feynman integrals

Pierre Vanhove



String Math 2017, Hamburg university, Hamburg  
based on [arXiv:1309.5865], [arXiv:1406.2664], [arXiv:1601.08181]

Spencer Bloch,



Matt Kerr



Scattering amplitudes are essential tools to understand a variety of physical phenomena from gauge theory to classical and quantum gravity

A convenient approach is to use modern unitarity methods for expanding the amplitude on a basis of integral functions

$$A^{\text{L-loop}} = \sum_{i \in \mathcal{B}(L)} \text{coeff}_i \text{Integral}_i + \text{Rational}$$

What are the intrinsic properties of amplitudes of QFT? How much can we understand about the amplitudes without having to compute them?

- ▶ What are the generic constraints on the integral coefficients?
- ▶ What are the elements of the basis of integral functions?

# Feynman Integrals: parametric representation

Any Feynman integrals with  $L$  loops and  $n$  propagators

$$I_\Gamma = \int \frac{\prod_{i=1}^L d^D \ell_i}{\prod_{i=1}^n d_i^{\nu_i}}$$

has the parametric representation

$$I_\Gamma = \Gamma(\nu - \frac{LD}{2}) \int_{x_i \geq 0} \frac{\mathcal{U}^{\nu - (L+1)\frac{D}{2}}}{(\mathcal{U} \sum_i m_i^2 x_i - \mathcal{V})^{\nu - L\frac{D}{2}}} \delta(x_n = 1) \prod_{i=1}^n \frac{dx_i}{x_i^{1-\nu_i}}$$


The Symanzik polynomials  $\mathcal{U}$  and  $\mathcal{V}$  are homogeneous in the  $x_1, \dots, x_n$

- ▶  $\mathcal{U}$  is of degree  $L$  in  $\mathbb{P}^{n-1}$
- ▶  $\mathcal{V}$  of degree  $L+1$  in  $\mathbb{P}^{n-1}$


# What are the Symanzik polynomials?

$$I_{\Gamma} = \Gamma\left(\nu - \frac{LD}{2}\right) \int_{x_i \geq 0} \frac{\mathcal{U}^{\nu - (L+1)\frac{D}{2}}}{(\mathcal{U} \sum_i m_i^2 x_i - \mathcal{V})^{\nu - L\frac{D}{2}}} \delta(x_n = 1) \prod_{i=1}^n \frac{dx_i}{x_i^{1-\nu_i}}$$


$\mathcal{U} = \det \Omega$  determinant of the period matrix of the graph  $\Omega_{ij} = \oint_{C_i} v_j$



(a)



(b)



(c)


$$\Omega_{2(a)} = \begin{pmatrix} x_1 + x_3 & x_3 \\ x_3 & x_2 + x_3 \end{pmatrix}; \quad \Omega_{3(b)} = \begin{pmatrix} x_1 + x_2 & x_2 & 0 \\ x_2 & x_2 + x_3 + x_5 + x_6 & x_3 \\ 0 & x_3 & x_3 + x_4 \end{pmatrix}$$

$$\Omega_{3(c)} = \begin{pmatrix} x_1 + x_4 + x_5 & x_5 & x_4 \\ x_5 & x_2 + x_5 + x_6 & x_6 \\ x_4 & x_6 & x_3 + x_4 + x_6 \end{pmatrix}$$

# What are the Symanzik polynomials?

$$I_{\Gamma} \propto \int_0^{\infty} \frac{\delta(1 - x_n)}{(\sum_i m_i^2 x_i - \mathcal{V}/\mathcal{U})^{n-L\frac{D}{2}}} \frac{1}{\mathcal{U}^{\frac{D}{2}}} \prod_{i=1}^n \frac{dx_i}{x_i^{1-\nu_i}}$$

$\mathcal{V}/\mathcal{U} = \sum_{1 \leq r \leq s \leq n} k_r \cdot k_s G(x_r/T_r, x_s/T_s; \Omega)$  sum of Green's function



$$G^{1-loop}(\alpha_r, \alpha_s; L) = -\frac{1}{2} |\alpha_s - \alpha_r| + \frac{1}{2} \frac{(\alpha_r - \alpha_s)^2}{T}.$$

# The geometry of a Feynman graph

The homogeneous polynomial of  $n$  variables and degree  $L + 1$  completely characterises the Feynman graph and its integral

$$\Phi_{\Gamma} = \mathcal{U} \times \left( \sum_{i=1}^n m_i^2 x_i \right) - \mathcal{V}$$

- ▶ We can recover both Symanzik polynomials
- ▶ Determines the graph topology
  - the number of propagators is the number of variables  $n$
  - the loop order is the degree minus one  $L = \deg(\Phi_{\Gamma}) - 1$
  - Number of vertices  $v = 1 + n - L$  from Euler characteristic


# From parametric representation to graph

The most general quadric polynomial in  $\mathbb{P}^2$

$$W_{2,3}(x_1, x_2, x_3) = \sum_{\substack{i_1+i_2+i_3=2 \\ i_r \geq 0}} w_{i_1, i_2, i_3} x_1^{i_1} x_2^{i_2} x_3^{i_3}$$

The graph has  $n = 3$  propagators,  $L = 1$  loop,  $v = 3$  vertices

This can only be a triangle graph



$$p_1 + p_2 + p_3 = 0; \quad p_i^2 \neq 0$$


$$\Phi_{\triangleright} = (x_1 + x_2 + x_3)(m_1^2 x_1 + m_2^2 x_2 + m_3^2 x_3) - (p_1^2 x_2 x_3 + p_2^2 x_1 x_3 + p_3^2 x_1 x_2)$$

# From parametric representation to graph

The most general cubic in  $\mathbb{P}^2$

$$W_{3,3} = \sum_{\substack{i_1+i_2+i_3=3 \\ i_r \geq 0}} w_{i_1, i_2, i_3} x_1^{i_1} x_2^{i_2} x_3^{i_3}$$

The graph has  $n = 3$  propagators,  $L = 2$  loops,  $v = 2$  vertices  
This can only be a sunset graph



$$\Phi_{\ominus} = (x_1 x_2 + x_1 x_3 + x_2 x_3)(m_1^2 x_1 + m_2^2 x_2 + m_3^2 x_3) - p^2 x_1 x_2 x_3$$



# From parametric representation to graph

The most general polynomial of degree  $n$  in  $\mathbb{P}^{n-1}$

$$W_{n,n} = \sum_{\substack{i_1 + \dots + i_n = n \\ i_r \geq 0}} w_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n}$$

The graph has  $n$  propagators,  $L = n - 1$  loops,  $v = 2$  vertices  
This can only be a  $n$ -loop sunset graphs



$$\Phi_n = \prod_{i=1}^n x_i \sum_{i=1}^n x_i^{-1} \sum_{i=1}^n m_i^2 x_i - p^2 \prod_{i=1}^n x_i$$

In general several graphs can occur in particular planar and non-planar topologies

# Feynman integral and periods

$$I_\Gamma = \Gamma\left(\nu - \frac{LD}{2}\right) \int_{\Delta_n} \Omega_\Gamma; \quad \Omega_\Gamma := \frac{\mathcal{U}^{\nu-(L+1)\frac{D}{2}}}{\Phi_\Gamma(x_i)^{\nu-L\frac{D}{2}}} \prod_{i=1}^{n-1} \frac{dx_i}{x_i^{1-\nu_i}}$$

$\Omega_\Gamma$  algebraic differential form on the complement of the graph hypersurface

$$\Omega_\Gamma \in H^{n-1}(\mathbb{P}^{n-1} \setminus X_\Gamma) \quad X_\Gamma := \{\Phi_\Gamma(x_i) = 0, x_i \in \mathbb{P}^{n-1}\}$$

The domain of integration is the simplex  $\Delta_n$

$$\Delta_n := \{x_1 \geq 0, \dots, x_n \geq 0 \mid [x_1, \dots, x_n] \in \mathbb{P}^{n-1}\}$$

# Feynman integral and periods

$$I_\Gamma = \Gamma\left(\nu - \frac{LD}{2}\right) \int_{\Delta_n} \Omega_\Gamma; \quad \Omega_\Gamma := \frac{\mathcal{U}^{\nu-(L+1)\frac{D}{2}}}{\Phi_\Gamma(x_i)^{\nu-L\frac{D}{2}}} \prod_{i=1}^{n-1} \frac{dx_i}{x_i^{\nu_i-1}}$$

The domain of integration is the simplex  $\Delta_n$

$$\Delta_n := \{x_1 \geq 0, \dots, x_n \geq 0 \mid [x_1, \dots, x_n] \in \mathbb{P}^{n-1}\}$$

with boundary contained in the normal crossings divisor

$$\partial\Delta_n \subset \mathbb{A}_n := \{x_1 \cdots x_n = 0\}$$


But  $\partial\Delta_n \cap X_\Gamma \neq \emptyset$  therefore  $\Delta_n \notin H_{n-1}(\mathbb{P}^{n-1} \setminus X_\Gamma)$

This is resolved by looking at the relative cohomology

$$H^\bullet(\mathbb{P}^{n-1} \setminus X_\Gamma; \mathbb{A}_n \setminus \mathbb{A}_n \cap X_\Gamma)$$

# Feynman integral and periods

$\mathcal{D}_n$  and  $X_\Gamma$  are separated by performing a series of iterated blowups of the complement of the graph hypersurface [Bloch, Esnault, Kreimer]



The Feynman integral are periods of the relative cohomology after performing the appropriate blow-ups

$$H^{n-1}(\widetilde{\mathbb{P}^{n-1}} \setminus \widetilde{X}_F; \widetilde{\mathcal{D}}_n \setminus \widetilde{\mathcal{D}}_n \cap \widetilde{X}_\Gamma)$$

# Feynman integral and periods

- ▶ In QFT one is interested in the  $\epsilon = (D - D_c)/2$  (e.g.  $D_c = 4$ ) expansion of the Feynman integral

$$I_\Gamma = \sum_{i \geq -n} c_i \epsilon^i$$

- ▶ The  $c_i$  are numerical periods [Belkale, Brosnan; Kontsevich, Zagier; Bogner, Weinzierl]

$$\mathfrak{M} := H^\bullet(\widetilde{\mathbb{P}^{n-1}} \setminus \widetilde{X}_F; \widetilde{\mathcal{D}}_n \setminus \widetilde{\mathcal{D}}_n \cap \widetilde{X}_\Gamma)$$

- ▶ The QFT questions: numbers of master integrals for amplitudes, their differential equations are now reformulated in a cohomological framework

# When physics and mathematics meet

The central questions about amplitudes in QFT can be reformulated as Riemann-Hilbert problem for periods

- ▶ Compute period explicitly



*Numerically or by series expansion in the physical region*

- ▶ Derive the local monodromy



*unitarity of the S-matrix*

- ▶ Construct a complete system of differential equations



*Relate this to the integration-by-part method used in QCD*

- ▶ Understand the new class of special functions that are needed



*What is needed beyond beyond elliptic multiple polylogarithm?*

# Differential equation

$$\mathfrak{M}(s_{ij}, m_i) := H^\bullet(\widetilde{\mathbb{P}^{n-1} \setminus X_F}; \widetilde{\mathcal{D}}_n \setminus \widetilde{\mathcal{D}}_n \cap \widetilde{X}_\Gamma)$$

Since  $\Omega_\Gamma$  varies when one changes the kinematic variables  $s_{ij}$  one needs to study a **variation of (mixed) Hodge structure**



Consequently the Feynman integral will satisfy a differential equation

$$L_{PF} I_\Gamma = S_\Gamma$$

The Picard-Fuchs operator will arise from the study of the variation of the differential in the cohomology when kinematic variables change

Generically there is an inhomogeneous term  $S_\Gamma \neq 0$

# The sunset family





# The sunset family

This talk will be focused on the special families of  $n$ -loop sunset graphs

$$\Phi_n = \prod_{i=1}^n x_i \sum_{i=1}^n x_i^{-1} \sum_{i=1}^n m_i^2 x_i - p^2 \prod_{i=1}^n x_i$$

- ▶ This family is a nice and important playground for understanding relations between Feynman integrals and periods
- ▶ This family leads to interesting motives : not mixed Tate, non trivial extensions
- ▶ Surprisingly rich: interesting Hodge structure, mirror symmetry
- ▶ For  $p^2 = m_1^2 = \dots = m_n^2$  [Broadhurst] found that special values of these sunset Feynman integrals are given by  $L$ -function evaluated in the critical band

# The sunset family

The graph polynomials  $\Phi_n = (-p^2) \prod_{i=1}^n x_i \times (1 - \frac{1}{p^2} \phi_n)$

$$\phi_n := \sum_{i=1}^n x_i^{-1} \times \sum_{i=1}^n m_i^2 x_i$$

- ▶  $\phi_n$  has a reflexive Newton polytope  $\Delta \subset \mathbb{R}^{n-1}$ .
- ▶ Its polar part  $\Delta^\circ$  has only integral points in  $\mathbb{R}^{n+1}$
- ▶  $\Delta^\circ$  is associated to a noncompact toric Fano  $n$ -fold  $\mathbb{P}_\Delta$

The sunset graphs lead to 1-parameter families of Calabi–Yau hypersurfaces in toric Fano  $n$ -folds

# The two-loop sunset integral

We consider the sunset integral in two Euclidean dimensions

$$\mathcal{I}_{\Theta}^2 = \int_{\Delta_3} \Omega_{\Theta}; \quad \Delta_3 := \{[x : y : z] \in \mathbb{P}^2 \mid x \geq 0, y \geq 0, z \geq 0\}$$

- ▶ The sunset integral is the integration of the 2-form

$$\Omega_{\Theta} = \frac{zdx \wedge dy + xdy \wedge dz + ydz \wedge dx}{(m_1^2 x + m_2^2 y + m_3^2 z)(xz + xy + yz) - p^2 xyz} \in H^2(\mathbb{P}^2 - \mathcal{E}_{p^2})$$

- ▶ The sunset family of open elliptic curve

$$\mathcal{E}_{p^2} = \{(m_1^2 x + m_2^2 y + m_3^2 z)(xz + xy + yz) - p^2 xyz = 0\}$$


- ▶ For  $m_1 = m_2 = m_3$  we have a modular curve  $\mathcal{E}_{p^2} \simeq X_1(6)$

# The differential operator: from the period

The analytic period of the elliptic curve around  $p^2 \sim \infty$  has the same integrand as the Feynman integral but we have just changed the domain of integration

$$\pi_0(p^2) := \int_{|x|=|y|=1} \Omega_\Theta$$

This is the imaginary part or the maximal cut of the amplitude


$$\Im(\mathcal{I}_\Theta(p^2)) = \oint_C \prod_{i=1}^3 \delta(\ell_i^2 - m_i^2) \delta(\ell_1 + \ell_2 + p) d^2 \ell_1 d^2 \ell_2$$

The other period is  $\pi_1(s) = \log(s) \pi_0(s) + \varpi_1(s)$  with  $\varpi_1(s)$  analytic is obtained by looking at different unitarity cut cutting less lines [Primo, Tancredi]

# The differential operator: from the period

The integral is the analytic period of the elliptic curve around  $p^2 \sim \infty$

$$\pi_0(p^2) := - \sum_{n \geq 0} \frac{1}{(p^2)^{n+1}} \left( \sum_{n_1+n_2+n_3=n} \left( \frac{n!}{n_1!n_2!n_3!} \right)^2 \prod_{i=1}^3 m_i^{2n_i} \right)$$

From the series expansion we can deduce the Picard-Fuch differential operator (the system has maximal unipotent monodromy [Lian, Todorov, Yau])

$$L_{\ominus} \pi_0(p^2) = 0$$

- ▶ With this method one easily derives the PF at all loop order for the all equal mass sunset and show the order(PF)=loop [Vanhove]
- ▶ Gives for the 3-loop sunset 2 unequal mass PF of order 4, and order 6 for 3 different masses [Vanhove; to appear]

# The differential equation

By general consideration we know that since the integrand is a top form we have

$$L_\Gamma I_\Gamma = \int_{\Delta_n} d\beta_\Gamma = - \int_{\partial\Delta_n} \beta_\Gamma = \mathcal{S}_\Gamma \neq 0$$

Writing the differential equation as  $\delta_s := s \frac{d}{ds}$   $\mathcal{S} = 1/p^2$

$$\left( \delta_s^2 + q_1(s)\delta_s + q_0(s) \right) \left( \frac{1}{s} I_\Theta(s) \right) = \mathcal{Y}_\Theta + \sum_{i=1}^3 \log(m_i^2) c_i(s)$$

# The differential equation

Using works from [del Angel, Müller-Stach] and [Doran, Kerr] we know that when rank of the  $D$ -module system of differential equations that  $y_\Theta$  is the Yukawa coupling


$$y_\Theta := \int_{\mathcal{E}(p^2)} \Omega_\Theta \wedge s \frac{d}{ds} \Omega_\Theta = \frac{2s^2 \prod_{i=1}^4 \mu_i - 4s \sum_i m_i^2 + 6}{\prod_{i=1}^4 (\mu_i^2 s - 1)}$$

The Yukawa coupling is the Wronskian of the Picard-Fuchs operator and only depends on the form of the Picard-Fuchs operator

$$y_\Theta = s \det \begin{pmatrix} \pi_0(s) & \pi_1(s) \\ \frac{d}{ds} \pi_0(s) & \frac{d}{ds} \pi_1(s) \end{pmatrix}$$

So far all we got can be deduced from the graph polynomial, and the associated Picard-Fuchs operator.

# The differential equation



$$(\delta_s^2 + q_1(s)\delta_s + q_0(s)) \left( \frac{1}{s} I_\Theta(s) \right) = y_\Theta + \sum_{i=1}^3 \log(m_i^2) c_i(s)$$

The mass dependent  $\log$ -terms come from derivative of partial elliptic integrals on globally well-defined algebraic 0-cycles arising from the punctures on the elliptic curve [Bloch, Kerr, Vanhove]

$$c_1(s) = \frac{d}{ds} \int_{q_2}^{q_3} \Omega_\Theta$$

They are rational function by construction.




# The 2-loop sunset integral as elliptic dilogarithm

The integral divided by a period of the elliptic curve is a function defined on the punctured torus [\[Bloch, Kerr, Vanhove\]](#)

$$\mathcal{I}_{\Theta} \equiv \frac{i\varpi_r}{\pi} \left( \mathcal{L}_2 \left\{ \frac{X}{Z}, \frac{Y}{Z} \right\} + \mathcal{L}_2 \left\{ \frac{Z}{X}, \frac{Y}{X} \right\} + \mathcal{L}_2 \left\{ \frac{X}{Y}, \frac{Z}{Y} \right\} \right) \mod \text{period}$$

- ▶  $\varpi_r$  is the elliptic curve period which is real on the line  $0 < p^2 < (m_1 + m_2 + m_3)^2$
- ▶ The sunset integral is the regulator period (with tame Milnor symbol) in the  $K_2$  of the elliptic curve [\[Bloch, Vanhove\]](#)

# The 2-loop sunset integral as elliptic dilogarithm



$E_s$

$$P_1 = [1, 0, 0];$$

$$Q_1 = [0, -m_3^2, m_2^2];$$

$$x(P_1)x(Q_1) = -1$$

$$P_2 = [0, 1, 0];$$

$$Q_2 = [-m_3^2, 0, m_1^2];$$

$$x(P_2)x(Q_2) = -1$$

$$P_3 = [0, 0, 1];$$

$$Q_3 = [-m_2^2, m_1^2, 0];$$

$$x(P_3)x(Q_3) = -1$$

Representing the ratio of the coordinates on the sunset cubic curve as functions on  $\mathcal{E}_\Theta \simeq \mathbb{C}^\times / q^\mathbb{Z}$

$$\frac{X}{Z}(x) = \frac{\theta_1(x/x(Q_1))\theta_1(x/x(P_3))}{\theta_1(x/x(P_1))\theta_1(x/x(Q_3))}$$

$$\frac{Y}{Z}(x) = \frac{\theta_1(x/x(Q_2))\theta_1(x/x(P_3))}{\theta_1(x/x(P_2))\theta_1(x/x(Q_3))}$$

$\theta_1(x)$  is the Jacobi theta function

$$\theta_1(x) = q^{\frac{1}{8}} \frac{x^{1/2} - x^{-1/2}}{i} \prod_{n \geq 1} (1 - q^n)(1 - q^n x)(1 - q^n/x).$$

# The 2-loop sunset integral as elliptic dilogarithm

$$\mathcal{L}_2 \left\{ \frac{X}{Z}, \frac{Y}{Z} \right\} = - \int_{x_0}^x \log \left( \frac{X}{Z}(y) \right) d \log y$$

Since

$$\begin{aligned} \int \log(\theta_1(x)) d \log x &= \sum_{n \geq 1} \int (\text{Li}_1(q^n x) + \text{Li}_1(q^n/x) + \text{cste}) d \log(x) \\ &= \sum_{n \geq 1} (\text{Li}_2(q^n x) - \text{Li}_2(q^n/x)) + \text{cste} \log(x) \end{aligned}$$

# The 2-loop sunset integral as elliptic dilogarithm

We find


$$\mathcal{J}_{\Theta}(s) \equiv \frac{i\omega_r}{\pi} \left( \hat{E}_2 \left( \frac{x(P_1)}{x(P_2)} \right) + \hat{E}_2 \left( \frac{x(P_2)}{x(P_3)} \right) + \hat{E}_2 \left( \frac{x(P_3)}{x(P_1)} \right) \right) \mod \text{periods}$$

where

$$\hat{E}_2(x) = \sum_{n \geq 0} (\text{Li}_2(q^n x) - \text{Li}_2(-q^n x)) - \sum_{n \geq 1} (\text{Li}_2(q^n/x) - \text{Li}_2(-q^n/x)) .$$

Close to the form given by [\[Brown, Levin\]](#). See as well [\[Adams, Bogner, Weinzierl\]](#)

# The three-loop sunset graph: integral



We look at the 3-loop sunset graph in  $D = 2$  dimensions

- ▶ The Feynman parametrisation is given by

$$I_{\oplus}^2(m_i; K^2) = \int_{x_i \geq 0} \frac{1}{(m_4^2 + \sum_{i=1}^3 m_i^2 x_i)(1 + \sum_{i=1}^3 x_i^{-1}) - K^2} \prod_{i=1}^3 \frac{dx_i}{x_i}$$

# three-loop sunset graph: differential equation

For the all equal mass case the geometry of the 3-loop sunset graph is a  $K3$  surface (Shioda-Inose family for  $\Gamma_1(6)^{+3}$ ) with Picard number 19 and discriminant of Picard lattice is 6

$$(m^2 + \sum_{i=1}^3 m^2 x_i)(1 + \sum_{i=1}^3 x_i^{-1}) \prod_{i=1}^3 x_i - p^2 \prod_{i=1}^3 x_i = 0$$

The  $t = p^2/m^2$  Picard-Fuchs equation

$$\begin{aligned} \left( t^2(t-4)(t-16) \frac{d^3}{dt^3} + 6t(t^2-15t+32) \frac{d^2}{dt^2} \right. \\ \left. + (7t^2-68t+64) \frac{d}{dt} + t-4 \right) J_{\oplus}^2(t) = -4! \end{aligned}$$

- One miracle is that this picard-fuchs operator is the symmetric square of the picard-fuchs operator for the sunset graph [Verrill]

# three-loop sunset graph: solution

- It is immediate to use the Wronskian method to solve the differential equation [Bloch, Kerr, Vanhove]


$$m^2 I_{\odot}^2(t) = 40\pi^2 \log(q) \varpi_1(\tau) \\ - 48\varpi_1(\tau) \left( 24\mathcal{L}i_3(\tau, \zeta_6) + 21\mathcal{L}i_3(\tau, \zeta_6^2) + 8\mathcal{L}i_3(\tau, \zeta_6^3) + 7\mathcal{L}i_3(\tau, 1) \right)$$

with  $\mathcal{L}i_3(\tau, z)$  [Zagier; Beilinson, Levin]


$$\mathcal{L}i_3(\tau, z) := \text{Li}_3(z) + \sum_{n \geq 1} (\text{Li}_3(q^n z) + \text{Li}_3(q^n z^{-1})) \\ - \left( -\frac{1}{12} \log(z)^3 + \frac{1}{24} \log(q) \log(z)^2 - \frac{1}{720} (\log(q))^3 \right).$$

- The 3-loop sunset integral is a regulator period of a motivic class of the  $K_3$  of the the  $K3$  surface [Bloch, Kerr, Vanhove]

# Mirror Symmetry



sunset



sunrise



# The sunset Gromov-Witten invariants

Around  $1/s = p^2 = \infty$  the sunset Feynman has the expansion

$$\mathcal{I}_{\ominus}(s) = -\pi_0 \left( 3R_0^3 + \sum_{\substack{\ell_1 + \ell_2 + \ell_3 = \ell > 0 \\ (\ell_1, \ell_2, \ell_3) \in \mathbb{N}^3 \setminus (0,0,0)}} \ell(1 - \ell R_0) N_{\ell_1, \ell_2, \ell_3} \prod_{i=1}^3 m_i^{2\ell_i} e^{\ell R_0} \right).$$

where the Kähler parameters are  $Q_i = m_i^2 e^{R_0}$  and  $R_0$  is the logarithmic Mahler measure defined by

$$R_0 := i\pi - \int_{|x|=|y|=1} \log(\Phi_{\ominus}(x, y)/(xy)) \frac{d \log x d \log y}{(2\pi i)^2}.$$

This is related to the holomorphic  $\pi_0(s)$  period near  $s = 1/p^2 = 0$

$$\pi_0 = s \frac{dR_0(s)}{ds}$$

# The sunset Gromov-Witten invariants

The numbers  $N_{\ell_1, \ell_2, \ell_3}$  are local Gromov-Witten expressed in terms of the virtual integer number of degree  $\ell$  rational curves by

$$N_{\ell_1, \ell_2, \ell_3} = \sum_{d|\ell_1, \ell_2, \ell_3} \frac{1}{d^3} n_{\frac{\ell_1}{d}, \frac{\ell_2}{d}, \frac{\ell_3}{d}}.$$

$\underline{\ell}$	(100)	$\overset{k>0}{(k00)}$	(110)	(210)	(111)	(310)	(220)	(211)	(221)
$N_{\underline{\ell}}$	2	$2/k^3$	-2	0	6	0	-1/4	-4	10
$n_{\underline{\ell}}$	2	0	-2	0	6	0	0	-4	10

$\underline{\ell}$	(410)	(320)	(311)	(510)	(420)	(411)	(330)	(321)	(222)
$N_{\underline{\ell}}$	0	0	0	0	0	0	-2/27	-1	-189/4
$n_{\underline{\ell}}$	0	0	0	0	0	0	0	-1	-48

# The sunset Gromov-Witten invariants

For the all equal masses case  $m_1 = m_2 = m_3 = 1$ , the mirror map is

$$Q = e^{R_0} = -q \prod_{n \geq 1} (1 - q^n)^{n\delta(n)}; \quad \delta(n) := (-1)^{n-1} \left( \frac{-3}{n} \right),$$

where  $\left( \frac{-3}{n} \right) = 0, 1, -1$  for  $n \equiv 0, 1, 2 \pmod{3}$ .

The local Gromov-Witten numbers

$$\begin{aligned} \frac{N_\ell}{6} = & 1, -\frac{7}{8}, \frac{28}{27}, -\frac{135}{64}, \frac{626}{125}, -\frac{751}{54}, \frac{14407}{343}, -\frac{69767}{512}, \frac{339013}{729}, -\frac{827191}{500}, \frac{8096474}{1331}, \\ & -\frac{367837}{16}, \frac{195328680}{2197}, -\frac{137447647}{392}, \frac{4746482528}{3375}, -\frac{23447146631}{4096}, \frac{115962310342}{4913}, \\ & -\frac{574107546859}{5832}, \frac{2844914597656}{6859}, -\frac{1410921149451}{800}, \frac{10003681368433}{1323}, \dots \end{aligned}$$

# The sunset mirror symmetry

- ▶ The sunset elliptic curve is embedded into a singular compactification  $X_0$  of the local Hori-Vafa 3-fold

$$Y := \{1 - s(m_1^2 x + m_2^2 y + m_3^2)(1 + x^{-1} + y^{-1}) + uv = 0\} \subset (\mathbb{C}^*)^2 \times \mathbb{C}^2,$$

limit of a family of elliptically-fibered CY 3-folds  $X_Z$

- ▶ The base given by  $\Phi_\Theta$  is a toric del Pezzo surface of degree 6
- ▶ We have an isomorphism of A- and B-model  $\mathbb{Z}$ -variation of Hodge structure

$$H^3(X_{z_0}) \cong H^{\text{even}}(X_{Q_0}^\circ),$$

and taking (the invariant part of) limiting mixed Hodge structure on both sides yields

the sunset Feynman integral given by the second regulator period of the motivic cohomology class is identified to the local Gromov-Witten prepotential for the 3-fold  $X$

# Mirror symmetry for elliptically fibered CY 3-fold

- In the degeneration limit the Yukawa coupling CY 3-fold  $X$  leads to the local Yukawa of the sunset elliptic curve

$$Y_{ijk} = \int_X \tilde{\Omega} \wedge \nabla_{\delta_i \delta_j \delta_k} \tilde{\Omega} \implies Y_{0ij}^{\text{loc}} \propto Y_{\Theta} = \int \Omega_{\Theta} \wedge \nabla_{\frac{d}{ds}} \Omega_{\Theta}$$

The holomorphic prepotential of [Huang, Klemm, Poretschkin]

$$F(Q_1, Q_2, Q_3, Q_4) = \frac{c_{ijk} t^i t^j t^k}{3!} + \frac{c_{ij}}{2!} t^i t^j + c_i t^i + c + \sum_{\beta \in H_2(M, \mathbb{Z})} n_0^{\beta} \text{Li}_3(Q^{\beta})$$

is mapped to the sunset integral with the identification of the Kähler parameter  $Q_r = \exp(2\pi i t_r) = m_r^2 Q$  for  $r = 1, 2, 3$  [Klemm private communication]

$$m_1^2 = \frac{(Q_1 Q_2 Q_4)^{\frac{1}{3}}}{Q_1^{\frac{2}{3}}}; m_2^2 = \frac{(Q_1 Q_2 Q_4)^{\frac{1}{3}}}{Q_2^{\frac{2}{3}}}; m_3^2 = \frac{(Q_1 Q_3 Q_4)^{\frac{1}{3}}}{Q_3^{\frac{2}{3}}}; Q = (Q_1 Q_2 Q_3 Q_4)^{\frac{1}{3}}$$

# Mirror symmetry for higher sunset integrals

- ☀ The same construction applies to the 3-loop sunset graph where  $\Phi_4 = 0$  defines a family of  $K3$
- ☀ The same is conjectured to be true for the 4-loop sunset graph where  $\Phi_5 = 0$  defines a family of CY 3-fold. [Doran, Kerr]
  - Not modular in general [Hulek, Verill].
  - Therefore (elliptic) polylogarithm not enough from 4-loop

At higher-loop loop the geometry is more intricate

- ☀ Need to go beyond the smoothness hypothesis for  $K_{\mathbb{P}^n_\Delta}$  used in [Lian, Todorov, Yau]
- ☀ Need to extend the construction of the motivic cohomology classes and the regulator period of [Doran, Kerr]

## The construction gives new way for computing amplitudes in QFT

- ▶ Efficient method for deriving Picard-Fuchs equation for Feynman integral in geometrical way
- ▶ Should help with the integration by part method and fix the ambiguities in the definition of the loop momentum

## Nice recent developments in mathematics

- ▶ [Deligne] conjectures and [Broadhurst] results on the relation between period and  $L$ -functions values (cf [Bloch, Kerr, Vanhove])
- ▶ Recent approach by [Zhou] using Hilbert transforms
- ▶ Relation to the Gamma class of [Golyushev, Zagier]